



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## FOREIGN AND INSULAR STATISTICAL REPORTS OF COUNTRIES AND CITIES—UNTABULATED.

AFRICA—*Kimberley*.—Month ended October 10, 1908. Estimated population, 54,141. Total number of deaths, 54, including 1 from enteric fever.

ARGENTINA—*Rosario de Santa Fe*.—Month of August, 1908. Estimated population, 154,663. Total number of deaths 267, including diphtheria 3, enteric fever 2, whooping cough 1, scarlet fever 4, and 30 from tuberculosis.

AUSTRIA—*Brünn*.—Months of August and September, 1908. Estimated population, 95,342. Total number of deaths 410, including diphtheria 3, enteric fever 2, scarlet fever 5, and 87 from tuberculosis.

BRAZIL—*Pernambuco*.—Two weeks ended September 30, 1908. Estimated population, 210,000. Total number of deaths, 286, including smallpox 38, measles 2, plague 2, leprosy 2, and 62 from tuberculosis.

CANADA—*British Columbia, Vancouver*.—Month ended October 31, 1908. Estimated population, 52,000. Total number of deaths, 51, including diphtheria 3, enteric fever 2, and 8 from tuberculosis. Cases: Diphtheria 15, enteric fever 19, scarlet fever 1, smallpox 2, and whooping cough 18.

Quebec—*Sherbrooke*.—Month of October, 1908. Estimated population, 14,709. Total number of deaths, 25, including enteric fever 1, and 4 from tuberculosis.

CUBA—*Sagua la Grande*.—Month ended October 30, 1908. Estimated population, 22,634. Total number of deaths, 35, including enteric fever 1, and 7 from tuberculosis.

FRANCE—*Calais*.—Month of October, 1908. Estimated population, 80,000. Total number of deaths, 141, including enteric fever 1, diphtheria 1, and 28 from tuberculosis.

*Cherbourg*.—Month of October, 1908. Estimated population, 43,948. Total number of deaths, 66, including enteric fever 7, diphtheria 1, and 13 from tuberculosis.

*Marseille*.—Month of October, 1908. Estimated population, 517,498. Total number of deaths, 770, including diphtheria 8, enteric fever 42, measles 5, whooping cough 1, and 103 from tuberculosis.

*St. Etienne*.—Two weeks ended October 31, 1908. Estimated population, 150,000. Total number of deaths, 132, including enteric fever 2 and 14 from tuberculosis. Cases: Enteric fever 10, diphtheria 1, and scarlet fever 6.

GREAT BRITAIN—*England and Wales*.—The deaths registered in 76 great towns in England and Wales during the week ended October 31, 1908, correspond to an annual rate of 15.3 per 1,000 population, which is estimated at 16,234,952.

*London*.—One thousand three hundred and ninety-six deaths were registered during the week, including measles 19, scarlet fever 13, diphtheria 16, enteric fever 11, whooping cough 7, tuberculosis 143, and 76 from diarrhea. The deaths from all causes correspond to an annual rate of 13.1 per 1,000. In Greater London 1,984 deaths were registered. In the "outer ring" the deaths included 10 from measles, 1 from scarlet fever, 8 from diphtheria, and 2 from whooping cough.

*Ireland*.—The average annual death rate represented by the deaths registered during the week ended October 31, 1908, in the 21 principal town districts of Ireland was 17 per 1,000 of the population, which is estimated at 1,131,959. The lowest rate was recorded in Lisburn, viz, 4.5, and the highest in Dundalk, viz, 35.9 per 1,000.

*Scotland*.—The deaths registered in 8 principal towns during the week ended October 31, 1908, correspond to an annual rate of 16.2 per 1,000 of the population, which is estimated at 1,839,038. The highest rate of mortality was recorded in Perth, viz, 22.2, and the lowest in Edinburgh, viz, 13 per 1,000. The aggregate number of deaths registered from all causes was 570, including enteric fever 2, diphtheria 11, scarlet fever 5, and 16 from whooping cough.

*HAWAII—Honolulu*.—Month of October, 1908. Estimated population, 39,306. Total number of deaths reported, 74, including enteric fever 3, diphtheria 1, and 12 from tuberculosis.

*JAMAICA—Kingston*.—Month of October, 1908. Estimated population, 52,065. Total number of deaths, 121, including enteric fever 9, and 26 from phthisis pulmonalis.

*MAURITIUS—Port Louis*.—Month of August, 1908. Estimated population, 376,635. Total number of deaths, 1,290, including enteric fever 4, measles 2, plague 8, and 78 from tuberculosis.

*NETHERLANDS—Schiedam*.—Month of October, 1908. Estimated population, 31,193. Total number of deaths, 30. Cause of deaths not given.

*SIBERIA—Vladivostok*.—Period from October 5 to 14, 1908. Estimated population, 79,699. Total number of deaths not reported. One death from typhus fever.

*SOCIETY ISLANDS—Tahiti*.—Five weeks ended October 31, 1908. Estimated population, 4,000. Total number of deaths 5, including enteric fever 1, and 2 from tuberculosis.

*URUGUAY—Montevideo*.—Month of August, 1908. Estimated population, 252,000. Total number of deaths, 441, including enteric fever 1, measles 1, scarlet fever 1, diphtheria 3, leprosy 1, and 53 from tuberculosis.